



# Parton Distributions and New Physics Searches: the Drell-Yan Forward-Backward Asymmetry as a Case Study

Juan Rojo, VU Amsterdam & Nikhef based on arXiv:2209.08115

**ATLAS PDF fitting Forum** 

14th October 2022, CERN/Zoom

### **BSM Searches with High-Mass Drell-Yan**

High-mass neutral-current Drell-Yan is a very sensitive processes for BSM searches

Resonant and EFT new physics can be probed with the invariant mass distributions, while offshell interference (e.g. Z' boson) one can use the Forward-Backward asymmetry



Surrent understanding of **large-x PDFs**: robust enough for present and future searches in DY?

At leading order, the triple differential cross-section in neutral-current Drell-Yan production is

$$\frac{\mathrm{d}^{3}\sigma}{\mathrm{d}m_{\ell\bar{\ell}}\,\mathrm{d}y_{\ell\bar{\ell}}\,\mathrm{d}\cos\theta^{*}} = \frac{\pi\alpha^{2}}{3m_{\ell\bar{\ell}}s} \left( (1+\cos^{2}(\theta^{*}))\sum_{q}S_{q}\left[f_{q}(x_{1},m_{\ell\bar{\ell}}^{2})f_{\bar{q}}(x_{2},m_{\ell\bar{\ell}}^{2}) + f_{q}(x_{2},m_{\ell\bar{\ell}}^{2})f_{\bar{q}}(x_{1},m_{\ell\bar{\ell}}^{2})\right] + \cos^{2}\theta^{*}\sum_{q}A_{q}\operatorname{sign}(y_{\ell\bar{\ell}})\left[f_{q}(x_{1},m_{\ell\bar{\ell}}^{2})f_{\bar{q}}(x_{2},m_{\ell\bar{\ell}}^{2}) - f_{q}(x_{2},m_{\ell\bar{\ell}}^{2})f_{\bar{q}}(x_{1},m_{\ell\bar{\ell}}^{2})\right] \right)$$

$$\overset{\text{invariant mass}}{\operatorname{soper angle}} \operatorname{Soper angle}$$

$$\begin{aligned} x_{1} &= \frac{m_{\ell\bar{\ell}}}{\sqrt{s}} \exp(y_{\ell\bar{\ell}}), \quad x_{2} = \frac{m_{\ell\bar{\ell}}}{\sqrt{s}} \exp(-y_{\ell\bar{\ell}}) \\ S_{q} &= e_{l}^{2} e_{q}^{2} + P_{\gamma Z} \cdot e_{l} v_{l} e_{q} v_{q} + P_{ZZ} \cdot (v_{l}^{2} + a_{l}^{2})(v_{q}^{2} + a_{q}^{2}) \\ A_{q} &= P_{\gamma Z} \cdot 2 e_{l} a_{l} e_{q} a_{q} + P_{ZZ} \cdot 8 v_{l} a_{l} v_{q} a_{q} , \\ P_{\gamma Z}(m_{\ell\bar{\ell}}) &= \frac{2m_{\ell\bar{\ell}}^{2}(m_{\ell\bar{\ell}}^{2} - m_{Z}^{2})}{\sin^{2}(\theta_{W}) \cos^{2}(\theta_{W}) \left[ (m_{\ell\bar{\ell}}^{2} - m_{Z}^{2})^{2} + \Gamma_{Z}^{2} m_{Z}^{2} \right]} \\ P_{ZZ}(m_{\ell\bar{\ell}}) &= \frac{m_{\ell\bar{\ell}}^{4}}{\sin^{4}(\theta_{W}) \cos^{4}(\theta_{W}) \left[ (m_{\ell\bar{\ell}}^{2} - m_{Z}^{2})^{2} + \Gamma_{Z}^{2} m_{Z}^{2} \right]} \end{aligned}$$

Express in terms of symmetric and antisymmetric parton luminosities



$$\mathcal{L}_{S,q}(m_{\ell\bar{\ell}}, y_{\ell\bar{\ell}}) \equiv f_q(x_1, m_{\ell\bar{\ell}}^2) f_{\bar{q}}(x_2, m_{\ell\bar{\ell}}^2) + f_q(x_2, m_{\ell\bar{\ell}}^2) f_{\bar{q}}(x_1, m_{\ell\bar{\ell}}^2) , \qquad \text{invariant under} \\ \mathcal{L}_{A,q}(m_{\ell\bar{\ell}}, y_{\ell\bar{\ell}}) \equiv \operatorname{sign}(y_{\ell\bar{\ell}}) \left[ f_q(x_1, m_{\ell\bar{\ell}}^2) f_{\bar{q}}(x_2, m_{\ell\bar{\ell}}^2) - f_q(x_2, m_{\ell\bar{\ell}}^2) f_{\bar{q}}(x_1, m_{\ell\bar{\ell}}^2) \right] \qquad x_1 \leftrightarrow x_2$$

Fractional Formattice Angle is defined in the hadronic CoM frame

$$\begin{aligned} \cos \theta^* &= \operatorname{sign}(y_{\ell \bar{\ell}}) \cos \theta \,, \\ \cos \theta &\equiv \frac{p_{\ell}^+ p_{\bar{\ell}}^- - p_{\ell}^- p_{\bar{\ell}}^+}{m_{\ell \bar{\ell}} \sqrt{m_{\ell \bar{\ell}}^2 + p_{\mathrm{T},\ell \bar{\ell}}^2}}, \quad p^{\pm} = p^0 \pm p^3 \end{aligned}$$

coincides with the lepton scattering angle in the partonic CoM frame

Express in terms of symmetric and antisymmetric parton luminosities



Secondination of symmetric and antisymmetric contributions in the Collins-Soper angle

A forward-backward (FB) asymmetry arises when antisymmetric lumi is non-zero

$$\frac{d^{3}\sigma}{dm_{\ell\bar{\ell}}dy_{\ell\bar{\ell}}d\cos(\theta^{*})}\Big|_{\mathrm{FB}} = \frac{d^{3}\sigma}{dm_{\ell\bar{\ell}}dy_{\ell\bar{\ell}}d\cos(\theta^{*})}\Big|_{\cos\theta^{*}} - \frac{d^{3}\sigma}{dm_{\ell\bar{\ell}}dy_{\ell\bar{\ell}}d\cos(\theta^{*})}\Big|_{-\cos\theta^{*}}$$
$$\frac{d^{3}\sigma}{dm_{\ell\bar{\ell}}dy_{\ell\bar{\ell}}d\cos(\theta^{*})}\Big|_{\mathrm{FB}} = \frac{2\pi\alpha^{2}\cos(\theta^{*})}{3m_{\ell\bar{\ell}}s}\sum_{q}A_{q}\mathscr{L}_{A,q}$$

At LO, properties of forward-backward asymmetry dictated by antisymmetric parton luminosity

One-dimensional distributions obtained from integrating the 3D ones

$$\begin{split} \frac{\mathrm{d}\sigma}{\mathrm{d}y_{\ell\bar{\ell}}} &= \int_{m_{\ell\bar{\ell}}^{\min}}^{\sqrt{s}} \mathrm{d}m_{\ell\bar{\ell}} \int_{-1}^{1} \mathrm{d}\cos\theta^* \frac{\mathrm{d}^3\sigma}{\mathrm{d}m_{\ell\bar{\ell}}\,\mathrm{d}y_{\ell\bar{\ell}}\,\mathrm{d}\cos\theta^*} \qquad \begin{array}{l} \text{Used frequently in} \\ global PDF fits \\ \end{array} \\ \frac{\mathrm{d}\sigma}{\mathrm{d}\cos\theta^*} &= \int_{m_{\ell\bar{\ell}}^{\min}}^{\sqrt{s}} \mathrm{d}m_{\ell\bar{\ell}} \int_{\mathrm{ln}(m_{\ell\bar{\ell}}/\sqrt{s})}^{\ln(\sqrt{s}/m_{\ell\bar{\ell}})} \mathrm{d}y_{\ell\bar{\ell}} \frac{\mathrm{d}^3\sigma}{\mathrm{d}m_{\ell\bar{\ell}}\,\mathrm{d}y_{\ell\bar{\ell}}\,\mathrm{d}\cos\theta^*} \qquad \begin{array}{l} \text{Used to extract EW} \\ parameters and also \\ proposed for PDF fits \\ \end{array} \\ \frac{\mathrm{d}\sigma}{\mathrm{d}\cos\theta^*} &= (1+\cos^2\theta^*) \sum_{q} g_{S,q} + \cos\theta^* \sum_{q} g_{A,q} \\ g_{A,q} &= \frac{\pi\alpha^2}{3s} \int_{m_{\ell\bar{\ell}}^{m}}^{\sqrt{s}} \frac{\mathrm{d}m_{\ell\bar{\ell}}}{m_{\ell\bar{\ell}}} A_q(m_{\ell\bar{\ell}}) \int_{\ln(m_{\ell\bar{\ell}}/\sqrt{s})}^{\ln(\sqrt{s}/m_{\ell\bar{\ell}})} \mathrm{d}y_{\ell\bar{\ell}}\,\mathcal{L}_{A,q}(m_{\ell\bar{\ell}},y_{\ell\bar{\ell}}) \end{split}$$

All PDF dependence into the **coefficients** g<sub>A,q</sub> and g<sub>,q</sub>, dependence in CS angle ``trivial"

#### Forward-Backward asymmetry

Free forward-backward asymmetry is defined to be proportional to the terms odd in CS angle:

$$A_{\rm fb}(\cos\theta^*) \equiv \frac{\frac{\mathrm{d}\sigma}{\mathrm{d}\cos\theta^*}(\cos\theta^*) - \frac{\mathrm{d}\sigma}{\mathrm{d}\cos\theta^*}(-\cos\theta^*)}{\frac{\mathrm{d}\sigma}{\mathrm{d}\cos\theta^*}(\cos\theta^*) + \frac{\mathrm{d}\sigma}{\mathrm{d}\cos\theta^*}(-\cos\theta^*)}, \quad \cos\theta^* > 0,$$

At LO the dependence on the CS angle **factorises** from the PDF dependence

$$A_{\rm fb}(\cos\theta^*) = \frac{\cos\theta^*}{(1+\cos^2(\theta^*))} \frac{\sum_q g_{A,q}}{\sum_{q'} g_{S,q'}}, \quad \cos\theta^* > 0$$

$$PDF\text{-independent} \quad PDF\text{-dependent}$$

Hence, to understand the high-mass behaviour of A<sub>FB</sub> it suffices to consider that of the antisymmetric PDF luminosities

$$g_{A,q} = rac{\pi lpha^2}{3s} \int\limits_{m_{\ell ar \ell}^{\min}}^{\sqrt{s}} rac{\mathrm{d}m_{\ell ar \ell}}{m_{\ell ar \ell}} A_q(m_{\ell ar \ell}) \int\limits_{\ln(m_{\ell ar \ell}/\sqrt{s})}^{\ln(\sqrt{s}/m_{\ell ar \ell})} \mathrm{d}y_{\ell ar \ell} \, \mathcal{L}_{A,q}(m_{\ell ar \ell}, y_{\ell ar \ell})$$

The following quantitative discussion is unaffected by higher-order QCD and EW corrections We will then carry out LHC phenomenology using NLO QCD+EW calculations

Free antisymmetric parton luminosities dictate the **behaviour of the FB asymmetry** 

Frey can also be expressed in terms of **sea-like** and **valence-like** PDF combinations

$$\begin{split} \mathcal{L}_{S,q}(m_{\ell\bar{\ell}}, y_{\ell\bar{\ell}}) &= \frac{1}{2} \left( f_q^+(x_1, m_{\ell\bar{\ell}}^2) f_q^+(x_2, m_{\ell\bar{\ell}}^2) - f_q^-(x_2, m_{\ell\bar{\ell}}^2) f_q^-(x_1, m_{\ell\bar{\ell}}^2) \right) \\ \mathcal{L}_{A,q}(m_{\ell\bar{\ell}}, y_{\ell\bar{\ell}}) &= \frac{\operatorname{sign}(y_{\ell\bar{\ell}})}{2} \left( f_q^-(x_1, m_{\ell\bar{\ell}}^2) f_q^+(x_2, m_{\ell\bar{\ell}}^2) - f_q^-(x_2, m_{\ell\bar{\ell}}^2) f_q^+(x_1, m_{\ell\bar{\ell}}^2) \right) \\ f_q^{\pm}(x, Q) &= f_q(x, Q) \pm f_{\bar{q}}(x, Q) \end{split}$$

In the **Z-peak region**, hierarchy of momentum fractions:  $x_1 \gg x_2$ 

$$\mathcal{L}_{A,u}(y_{\ell ar{\ell}}, m_{\ell ar{\ell}}) pprox rac{1}{2} f_u^-(x_1, m_{\ell ar{\ell}}^2) f_u^+(x_2, m_{\ell ar{\ell}}^2) \qquad x_1 = rac{m_{\ell ar{\ell}}}{\sqrt{s}} \exp(y_{\ell ar{\ell}}), \quad x_2 = rac{m_{\ell ar{\ell}}}{\sqrt{s}} \exp(-y_{\ell ar{\ell}})$$

Hence measurements of the FB asymmetry on the Z-peak can constrain valence quark PDFs in addition, also the weak mixing angle can be measured

The integral over valence quark PDFs is a positive quantity (sum rules), it then follows that the forward-backward asymmetry is **positive-definite in the Standard Model?** 

very attractive feature for robust BSM searches!

Sector AFB has been shown to be particularly constraining for quark and down valence PDFs



 $\mathcal{L}_{A,u}(y_{\ell\bar{\ell}}, m_{\ell\bar{\ell}}) \approx \frac{1}{2} f_u^-(x_1, m_{\ell\bar{\ell}}^2) f_u^+(x_2, m_{\ell\bar{\ell}}^2)$ 

Most studies of AFB focus on Z-peak region What about higher invariant masses?

### **Antisymmetric PDF luminosities**

Search At the Z-peak region we probe valence PDFs: antisymmetric parton lumis behave valence-like



### **Antisymmetric PDF luminosities**

Search At the Z-peak region we probe valence PDFs: antisymmetric parton lumis behave valence-like



Same as we go to **2 TeV**, and note good agreement between PDF fits



### **Antisymmetric PDF luminosities**

Sector At 5 TeV, very different behaviour in NNPDF4.0: AFB may not be positive definite after all



Same as we go to 2 TeV, and note good agreement between PDF fits



### Symmetric PDF luminosities

On the other hand, symmetric parton luminosities are in good qualitative agreement even at very high masses, with NNPDF4.0 displaying the largest PDF uncertainties



How can this behaviour be explained?

### Positive or negative asymmetry?

Antisymmetric luminosity depends on relative rate of decrease of the quark and antiquark PDFs



AFB sensitive to **subtle PDF property**: difference in decrease rates of large-*x* quarks vs antiquarks

Quantified by the effective asymptotic exponents, which illustrate richer structure in NNPDF4.0



### Positive or negative asymmetry?

Evaluate the **PDF-dependent coefficient** of the forward-backward asymmetry in LO QCD:



Prediction: for NNPDF4.0 the **forward-backward asymmetry eventually vanishes** at high masses, for the other groups the predictions are mass-independent

# LHC phenomenology

Validate our LO interpretation with realistic LHC simulations based on mg5\_aMC with NLO QCD and EW corrections and with same fiducial selection cuts as in the ATLAS/CMS measurements



As well known, clearly positive FB asymmetry with good agreement between PDF fits

What happens at higher dilepton masses?

# LHC phenomenology

For dilepton masses > 3 TeV, same qualitative behaviour, with clearly positive AFB



However, we know from the LO analysis that extrapolation to yet high masses may change the qualitative behaviour

# LHC phenomenology

For dilepton masses > 5 TeV, AFB vanishes for NNPDF4.0, while other groups extrapolate



PDF uncertainties differ between PDF groups, with NNPDF4.0 displaying the largest ones

# Summary

As opposed to common lore, the forward-backward asymmetry in neutral-current Drell-Yan is not positive-definite in the Standard Model

Explained by the behaviour of the antisymmetric parton luminosities, which probe features of the large-x PDFs not accessible with processes such as DY rapidity distributions

Extrapolation to the large-x region of PDFs (and uncertainties) depends on methodological assumptions carried by PDF groups and on how much large-x data considered

Our findings emphasise that a careful understanding of large-x PDFs is crucial in order to robustly search for BSM physics in the high-mass region ...

sections) measurements at high mass provide **unique sensitivity to large-***x* **PDFs** 

Motivates revisiting other high-mass processes sensitive to both PDFs and BSM, from jets to top quark pair production